Product Datasheet

Technical Data

Formula						
Molecular Weight	N/A	**CAS No.**	11078-21-0			
Solubility (25°C)*	In vitro	DMSO	100 mg/mL			
		Ethanol	8 mg/mL			
		Water	Insoluble			
	In vivo (Add solvents to the product individually and in order)	Oral Administration	Homogeneous suspension	CMC-NA	≥5mg/ml	Taking the 1 mL working solution as an example, add 5 mg of this product to 1 ml of CMC-Na solution, mix evenly to obtain a homogeneous suspension with a final concentration of 5 mg/ml.
		Injection	Clear solution	5%DMSO [1]→ 40%PEG300 [2]→ 5%Tween80 [3]→ 50%ddH2O Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	1.650mg/ml	Taking the 1 mL working solution as an example, add 50 μL of 33 mg/ml clarified DMSO stock solution to 400 μL of PEG300, mix evenly to clarify it; add 50 μL of Tween80 to the above system, mix evenly to clarify; then continue to add 500 μL of ddH2O to adjust the volume to 1 mL. The mixed solution should be used immediately for optimal results.
			Clear solution	5% DMSO [1]→ 95% Corn oil Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	0.600mg/ml	Taking the 1 mL working solution as an example, add 50 μL of 12 mg/ml clear DMSO stock solution to 950 μL of corn oil and mix evenly. The mixed solution should be used immediately for optimal results.

* <1 mg/ml means slightly soluble or insoluble.
* Please note that Capsid Bio tests the solubility of all compounds in-house, and the actual solubility may differ slightly from published values. This is normal and is due to slight batch-to-batch variations.
* Room temperature shipping (Stability testing shows this product can be shipped at 2-8°C for storage purposes.)

Preparing Stock Solutions

Biological Activity

Description	Filipin complex is a potent polyene macrolide antifungal antibiotic that integrates into membranes, sequestering cholesterol into complexes. Its treatment reduces **PRRSV** genome copies by 61% and lowers the PRRSV titer. It also acts as a fluorescent histochemical stain prone to autooxidation, commonly used in diagnosing Niemann-Pick Disease Type C (NP-C), a neurodegenerative lysosomal storage disorder. Researchers also employ it to study unesterified cholesterol distribution in cell and animal models of neurodegenerative diseases like NP-C and Sanfilippo syndrome (MPS IIIA).

References

https://pubmed.ncbi.nlm.nih.gov/21986526/ https://pubmed.ncbi.nlm.nih.gov/38112248/

RETURN POLICY

Capsid Bio's Unconditional Return Policy ensures a smooth online shopping experience for our customers. If you are in any way unsatisfied with your purchase, you may return any item(s) within 7 days of receiving it. In the event of product quality issues, either protocol related or product related problems, you may return any item(s) within 365 days from the original purchase date. Please follow the instructions below when returning products.

SHIPPING & STORAGE

Capsid Bio products are transported at room temperature. If you receive the product at room temperature, please rest assured, the Capsid Bio Quality Inspection Department has conducted experiments to verify that the normal temperature placement of one month will not affect the biological activity of powder products. After collecting, please store the product according to the requirements described in the datasheet. Most Capsid Bio products are stable under the recommended conditions.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

EU Headquarters
Gajeva ulica 28, HR-10000 Zagreb, Croatia

US Headquarters
12222 Merit Dr STE 130 Dallas, Texas, 75251

UK Headquarters
121 ABILITY HOUSE, BROOKER ROAD, EN9 1JH, ESSEX

Orders: Sales@capsidbio.com
Enquiries: info@capsidbio.com
www.capsidbio.com