Product Datasheet

Biological Description

Specificity	Phospho-p27 KIP 1 (Ser10) Antibody [D8F10] detects endogenous levels of p27 KIP 1 only when phosphorylated at Ser10.
Background	P27Kip1 is a cyclin-dependent kinase inhibitor (CKI) that regulates cell cycle progression during the G1 phase by interacting with and inhibiting cyclin-dependent kinase (CDK)/cyclin complexes. Regulation of p27 expression and function involves transcriptional and translational mechanisms, as well as degradation by the 26S proteasome and miRNA. Serine 10 (S10) is the major phosphorylation site on p27Kip1. Phosphorylation of p27Kip1 at serine 10 (pS10-p27Kip1) influences its stability, cellular localization, and interactions with CDKs. Nuclear levels of pS10-p27Kip1 vary during the cell cycle, peaking in the G0 and G1 phases and decreasing in the S phase. pS10-p27Kip1 primarily interacts with CDK2, CDK4, and CDK6, inhibiting CDK2 activity, especially when associated with cyclin E. The stability of pS10-p27Kip1 is affected by its interaction with CDKs, indicating its role in regulating cell cycle progression. Decreased levels of pS10-p27Kip1 in the S phase are due to increased nuclear exit, facilitating cell cycle progression. S10 phosphorylation is necessary for T157 phosphorylation by AKT and subsequent protein localization in the cytoplasm.

Usage Information

Application	WB, IP, IHC	Dilution	WB: 1:500-1:20000; IP: 1:50; IHC: 1:100
Reactivity	Human, Mouse, Rat, Monkey		
Source	Rabbit Monoclonal Antibody	MW	22 kDa
Storage Buffer	PBS, pH 7.2+50% Glycerol+0.05% BSA+0.01% NaN_3		
Storage (from the date of receipt)	–20°C (avoid freeze-thaw cycles), 2 years		

Experimental Methods

WB

Experimental Protocol:

Sample preparation
1. Tissue: Lyse the tissue sample by adding an appropriate volume of ice-cold **RIPA/NP-40 Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**，and homogenize the tissue at a low temperature or lyse it by sonication on ice, then incubate on ice for 30 minutes.
2. Adherent cell: Aspirate the culture medium and transfer the cells into an EP tube. Wash the cells with ice-cold PBS twice. Add an appropriate volume of **RIPA/NP-40 Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**, sonicate to lyse the cells, and incubate on ice for 30 minutes.
3. Suspension cell: Transfer the culture medium to a pre-cooled centrifuge tube. Centrifuge and aspirate the supernatant. Wash the cells with ice-cold PBS twice.Add an appropriate volume of **RIPA/NP-40 Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**, sonicate to lyse the cells, and incubate on ice for 30 minutes.
4. Place the lysate into a pre-cooled microcentrifuge tube. Centrifuge at 4°C for 15 min. Collect the supernatant;
5. Remove a small volume of lysate to determine the protein concentration;
6. Combine the lysate with protein loading buffer. Boil 20 µL sample under 95-100°C for 5 min. Centrifuge for 5 min after cool down on ice.

Electrophoretic separation
1. According to the concentration of extracted protein, load appropriate amount of protein sample and marker onto SDS-PAGE gels for electrophoresis. Recommended separating gel (lower gel) concentration: 10%. Reference Table for Selecting SDS-PAGE Separation Gel Concentrations
2. Power up 80V for 30 minutes. Then the power supply is adjusted (110 V~150 V), the Marker is observed, and the electrophoresis can be stopped when the indicator band of the predyed protein Marker where the protein is located is properly separated. (Note that the current should not be too large when electrophoresis, too large current (more than 150 mA) will cause the temperature to rise, affecting the result of running glue. If high currents cannot be avoided, an ice bath can be used to cool the bath.)

Transfer membrane
1. Take out the converter, soak the clip and consumables in the pre-cooled converter;
2. Activate PVDF membrane with methanol for 1 min and rinse with transfer buffer;
3. Install it in the order of "black edge of clip - sponge - filter paper - filter paper - glue -PVDF membrane - filter paper - filter paper - sponge - white edge of clip";
4. The protein was electrotransferred to PVDF membrane. (0.45 µm PVDF membrane is recommended) Reference Table for Selecting PVDF Membrane Pore Size Specifications
Recommended conditions for wet transfer: **200 mA, 60 min**.
(Note that the transfer conditions can be adjusted according to the protein size. For high-molecular-weight proteins, a higher current and longer transfer time are recommended. However, ensure that the transfer tank remains at a low temperature to prevent gel melting.)

Block
1. After electrotransfer, wash the film with TBST at room temperature for 5 minutes;
2. Incubate the film in the blocking solution **(recommending 5% BSA solution)** for 1 hour at room temperature;
3. Wash the film with TBST for 3 times, 5 minutes each time.

Antibody incubation
1. Use 5% skim milk powder to prepare the primary antibody working liquid (recommended dilution ratio for primary antibody 1:1000), gently shake and incubate with the film at 4°C overnight;
2. Wash the film with TBST 3 times, 5 minutes each time;
3. Add the secondary antibody to the blocking solution and incubate with the film gently at room temperature for 1 hour;
4. After incubation, wash the film with TBST 3 times for 5 minutes each time.

Antibody staining
497. Add the prepared ECL luminescent substrate (or select other color developing substrate according to the second antibody) and mix evenly;
2. Incubate with the film for 1 minute, remove excess substrate (keep the film moist), wrap with plastic film, and expose in the imaging system.

IF

Experimental Protocol：

Specimen Preparation
1. Aspirate liquid, then cover cells to a depth of 2–3 mm with 4% Paraformaldehyde diluted in 1X PBS.
NOTE: Paraformaldehyde is toxic, use only in a fume hood.
2. Fix cells for 15 min at room temperature.
3. Aspirate fixative, rinse three times in 1X PBS for 5 min each.
4. Proceed with Immunostaining.

Immunostaining

Experimental Methods

	1. Add theblocking buffer and incubate for 60 min at RT. 2. Prepare primary antibody diluent in antibody dilution buffer as recommended . 3. Aspirate blocking solution, apply diluted primary antibody. 4. Incubate overnight at 4°C. 5. Rinse three times in 1X PBS for 5 min each. 6. Incubate specimens in fluorochrome-conjugated secondary antibody diluted in antibody dilution buffer for 1–2 hr at room temperature in the dark. 7. Rinse three times in 1X PBS for 5 min each. 8. Mount slides usingmounting medium with DAPI and cover with coverslips. 9. For best results, allow mountant to cure overnight at room temperature. For long-term storage, store slides flat at 4°C protected from light.
IHC	Experimental Protocol： Deparaffinization/Rehydration 1. Deparaffinize/hydrate sections: 2. Incubate sections in three washes of xylene for 5 min each. 3. Incubate sections in two washes of 100% ethanol for 10 min each. 4. Incubate sections in two washes of 95% ethanol for 10 min each. 5. Wash sections two times in dH2O for 5 min each. 6.Antigen retrieval: For Citrate: Heat slides in a microwave submersed in 1X citrate unmasking solution until boiling is initiated; continue with 10 min at a sub-boiling temperature (95°-98°C). Cool slides on bench top for 30 min. Staining 1. Wash sections in dH2O three times for 5 min each. 2. Incubate sections in 3% hydrogen peroxide for 10 min. 3. Wash sections in dH2O two times for 5 min each. 4. Wash sections in wash buffer for 5 min. 5. Block each section with 100–400 µl of blocking solution for 1 hr at room temperature. 6. Remove blocking solution and add 100–400 µl primary antibody diluent in to each section. Incubate overnight at 4°C. 7. Remove antibody solution and wash sections with wash buffer three times for 5 min each. 8. Cover section with 1–3 drops HRPas needed. Incubate in a humidified chamber for 30 min at room temperature. 9. Wash sections three times with wash buffer for 5 min each. 10. Add DAB Chromogen Concentrate to DAB Diluent and mix well before use. 11. Apply 100–400 µl DAB to each section and monitor closely. 1–10 min generally provides an acceptable staining intensity. 12. Immerse slides in dH2O. 13. If desired, counterstain sections with hematoxylin. 14. Wash sections in dH2O two times for 5 min each. 15. Dehydrate sections: Incubate sections in 95% ethanol two times for 10 sec each; Repeat in 100% ethanol, incubating sections two times for 10 sec each; Repeat in xylene, incubating sections two times for 10 sec each. 16. Mount sections with coverslips and mounting medium.


References

https://pubmed.ncbi.nlm.nih.gov/25483085/

Application Data

IHC

Validated by Capsid Bio



Immunohistochemical analysis of formalin fixed paraffin embedded human prostate cancer tissue with F1167 at 1/100 dilution.

WB

Validated by Capsid Bio



Lane 1: HeLa
Lane 2: HeLa (Alkaline Phosphatase,1h)
Lane 3: NIH/3T3
Lane 4: NIH/3T3 (Alkaline Phosphatase,1h)
Lane 5: C6
Lane 6: C6 (Alkaline Phosphatase,1h)