Product Datasheet

Technical Data

Formula	$C_{20}H_{28}O$					
Molecular Weight	284.44	**CAS No.**	116-31-4			
Solubility (25°C)*	In vitro	DMSO	57 mg/mL (200.39 mM)			
		Ethanol	57 mg/mL (200.39 mM)			
		Water	Insoluble			
	In vivo (Add solvents to the product individually and in order)	Oral Administration	Homogeneous suspension	CMC-NA	≥5mg/ml	Taking the 1 mL working solution as an example, add 5 mg of this product to 1 ml of CMC-Na solution, mix evenly to obtain a homogeneous suspension with a final concentration of 5 mg/ml.
		Injection	Clear solution	5%DMSO [1]→ 40%PEG300 [2]→ 5%Tween80 [3]→ 50%ddH2O Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	2.850mg/ml (10.02mM)	Taking the 1 mL working solution as an example, add 50 μL of 57 mg/ml clarified DMSO stock solution to 400 μL of PEG300, mix evenly to clarify it; add 50 μL of Tween80 to the above system, mix evenly to clarify; then continue to add 500 μL of ddH2O to adjust the volume to 1 mL. The mixed solution should be used immediately for optimal results.
			Clear solution	5% DMSO [1]→ 95% Corn oil Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	1.400mg/ml (4.92mM)	Taking the 1 mL working solution as an example, add 50 μL of 28 mg/ml clear DMSO stock solution to 950 μL of corn oil and mix evenly. The mixed solution should be used immediately for optimal results.

* <1 mg/ml means slightly soluble or insoluble.
* Please note that Capsid Bio tests the solubility of all compounds in-house, and the actual solubility may differ slightly from published values. This is normal and is due to slight batch-to-batch variations.
* Room temperature shipping (Stability testing shows this product can be shipped at 2-8°C for storage purposes.)

Preparing Stock Solutions

Concentration \ Volume \ Mass	1 mg	5 mg	10 mg
1 mM	3.5157 mL	17.5784 mL	35.1568 mL
5 mM	0.7031 mL	3.5157 mL	7.0314 mL
10 mM	0.3516 mL	1.7578 mL	3.5157 mL
50 mM	0.0703 mL	0.3516 mL	0.7031 mL

Biological Activity

Description	All trans-Retinal (Retinaldehyde, Vitamin A aldehyde, Retinene), a component of the retinoid cycle, is converted to retinoic acid in vivo by the action of retinal dehydrogenase. Retinoic acid is a ligand for both RAR and RXR. All trans-retinal ias a potent photosensitizer.

RETURN POLICY

Capsid Bio's Unconditional Return Policy ensures a smooth online shopping experience for our customers. If you are in any way unsatisfied with your purchase, you may return any item(s) within 7 days of receiving it. In the event of product quality issues, either protocol related or product related problems, you may return any item(s) within 365 days from the original purchase date. Please follow the instructions below when returning products.

SHIPPING & STORAGE

Capsid Bio products are transported at room temperature. If you receive the product at room temperature, please rest assured, the Capsid Bio Quality Inspection Department has conducted experiments to verify that the normal temperature placement of one month will not affect the biological activity of powder products. After collecting, please store the product according to the requirements described in the datasheet. Most Capsid Bio products are stable under the recommended conditions.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

EU Headquarters
Gajeva ulica 28, HR-10000 Zagreb, Croatia

US Headquarters
12222 Merit Dr STE 130 Dallas, Texas, 75251

UK Headquarters
121 ABILITY HOUSE, BROOKER ROAD, EN9 1JH, ESSEX

Orders: Sales@capsidbio.com
Enquiries: info@capsidbio.com
www.capsidbio.com