Product Datasheet

Technical Data

Formula	$C_{34}H_{24}N_6Na_4O_{14}S_4$					
Molecular Weight	960.81	**CAS No.**	72-57-1			
Solubility (25°C)*	In vitro	DMSO	100 mg/mL (104.07 mM)			
		Water	100 mg/mL (104.07 mM)			
		Ethanol	Insoluble			
	In vivo (Add solvents to the product individually and in order)	Oral Administration	Homogeneous suspension	CMC-NA	≥5mg/ml	Taking the 1 mL working solution as an example, add 5 mg of this product to 1 ml of CMC-Na solution, mix evenly to obtain a homogeneous suspension with a final concentration of 5 mg/ml.

* <1 mg/ml means slightly soluble or insoluble.
* Please note that Capsid Bio tests the solubility of all compounds in-house, and the actual solubility may differ slightly from published values. This is normal and is due to slight batch-to-batch variations.
* Room temperature shipping (Stability testing shows this product can be shipped at 2-8°C for storage purposes.)

Preparing Stock Solutions

Biological Activity

Description	Trypan Blue (Visionblue, Diphenyl Blue, Direct Blue 14) is a colorimetric dye that stains dead cells with a blue color easily observed using light microscopy at low resolution.
In vitro	1.1 Preparation of the stock solution Dissolve 400 mg of Trypan Blue in 100 mL 0.85% Nacl to obtain 0.4% of this compound. Note: It is recommended to store the stock solution at -20℃ -80℃ away from light and avoid repetitive freeze-thaw cycles. 1.2 Preparation of this compound working solution Dilute the stock solution in 0.85% Nacl to obtain 0.04% of this chemical working solution. Note: Please adjust the concentration of this chemical working solution according to the actual situation. Cell staining 2.1 For suspension cells, centrifuge at 1000 g at 4℃ for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. For adherent cells, discard the cell culture medium, and add trypsin to dissociate cells to make a single-cell suspension. Centrifuge at 1000 g at 4℃ for 3-5 minutes and then discard the supernatant. Wash twice with PBS, 5 minutes each time. 2.2 Add 1 mL of this chemical working solution, and then incubate at room temperature for 5 minutes. 2.3 Centrifuge at 400 g at 4℃ for 3-4 minutes and then discard the supernatant. 2.4 Wash twice with PBS, 5 minutes each time. 2.5 Resuspend cells with serum-free cell culture medium or PBS, and then detect by fluorescence microscope.

References

https://pubmed.ncbi.nlm.nih.gov/27371594/

RETURN POLICY

Capsid Bio's Unconditional Return Policy ensures a smooth online shopping experience for our customers. If you are in any way unsatisfied with your purchase, you may return any item(s) within 7 days of receiving it. In the event of product quality issues, either protocol related or product related problems, you may return any item(s) within 365 days from the original purchase date. Please follow the instructions below when returning products.

SHIPPING & STORAGE

Capsid Bio products are transported at room temperature. If you receive the product at room temperature, please rest assured, the Capsid Bio Quality Inspection Department has conducted experiments to verify that the normal temperature placement of one month will not affect the biological activity of powder products. After collecting, please store the product according to the requirements described in the datasheet. Most Capsid Bio products are stable under the recommended conditions.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

EU Headquarters
Gajeva ulica 28, HR-10000 Zagreb, Croatia

US Headquarters
12222 Merit Dr STE 130 Dallas, Texas, 75251

UK Headquarters
121 ABILITY HOUSE, BROOKER ROAD, EN9 1JH, ESSEX

Orders: Sales@capsidbio.com
Enquiries: info@capsidbio.com
www.capsidbio.com