Product Datasheet

Technical Data

Formula	$C_{43}H_{69}NO_{12}$					
Molecular Weight	792.01	**CAS No.**	104987-12-4			
Solubility (25°C)*	In vitro	DMSO	100 mg/mL (126.26 mM)			
		Ethanol	100 mg/mL (126.26 mM)			
		Water	Insoluble			
	In vivo (Add solvents to the product individually and in order)	Oral Administration	Homogeneous suspension	CMC-NA	≥5mg/ml	Taking the 1 mL working solution as an example, add 5 mg of this product to 1 ml of CMC-Na solution, mix evenly to obtain a homogeneous suspension with a final concentration of 5 mg/ml.
		Injection	Clear solution	5%DMSO [1] → 40%PEG300 [2] → 5%Tween80 [3] → 50%ddH2O Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	5.000mg/ml (6.31mM)	Taking the 1 mL working solution as an example, add 50 μL of 100 mg/ml clarified DMSO stock solution to 400 μL of PEG300, mix evenly to clarify it; add 50 μL of Tween80 to the above system, mix evenly to clarify; then continue to add 500 μL of ddH2O to adjust the volume to 1 mL. The mixed solution should be used immediately for optimal results.
			Clear solution	5% DMSO [1] → 95% Corn oil Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	1.650mg/ml (2.08mM)	Taking the 1 mL working solution as an example, add 50 μL of 33 mg/ml clear DMSO stock solution to 950 μL of corn oil and mix evenly. The mixed solution should be used immediately for optimal results.

* <1 mg/ml means slightly soluble or insoluble.
* Please note that Capsid Bio tests the solubility of all compounds in-house, and the actual solubility may differ slightly from published values. This is normal and is due to slight batch-to-batch variations.
* Room temperature shipping (Stability testing shows this product can be shipped at 2-8°C for storage purposes.)

Preparing Stock Solutions

Concentration \ Volume / Mass	1 mg	5 mg	10 mg
1 mM	1.2626 mL	6.3131 mL	12.6261 mL
5 mM	0.2525 mL	1.2626 mL	2.5252 mL
10 mM	0.1263 mL	0.6313 mL	1.2626 mL
50 mM	0.0253 mL	0.1263 mL	0.2525 mL

Biological Activity

Description	Ascomycin (FK520, FR 900520, Immunomycin), an FK-506 analog, is a neutral macrolide **immunosuppressant** that prevents rejection after an organ transplant. Phase 3.
Targets	calcineurin phosphatase [2] 49 nM
In vitro	Ascomycin (FK520), isolated from Streptomyces, suppresses immune response in vitro with IC50 of 0.55 nM for mouse mixed lymphocyte. [1] It inhibits calcineurin phosphatase with IC50 of 49 nM through formation of an FKBP12-FK520-calcineurin ternary complex. [2] This compound also accelerates the rate of nerve regeneration, and promotes neurite outgrowth. [2] It shows antimalarial effects via inhibition of the chaperone activity of this bifunctional protein. [3]
In vivo	Ascomycin (FK520) (3.2 mg/kg, i.m.) clearly prolongs skin allograft survival in rats. [1] When infused into the rat hippocampus at 50 or 100 μM, this compound produces anticonvulsant effect against picrotoxin-induced seizures. [4]

Protocol (from reference)

Animal Study: [1]

- **Animal Models**
  Recipient WKA rats transplanted with F344 skin allografts.
- **Dosages**
  ~32 mg/kg 5 days a week.
- **Administration**
  i.m.

References

https://pubmed.ncbi.nlm.nih.gov/2461926/ https://pubmed.ncbi.nlm.nih.gov/12183690/ https://pubmed.ncbi.nlm.nih.gov/15776382/

+ Expand

RETURN POLICY

Capsid Bio's Unconditional Return Policy ensures a smooth online shopping experience for our customers. If you are in any way unsatisfied with your purchase, you may return any item(s) within 7 days of receiving it. In the event of product quality issues, either protocol related or product related problems, you may return any item(s) within 365 days from the original purchase date. Please follow the instructions below when returning products.

SHIPPING & STORAGE

Capsid Bio products are transported at room temperature. If you receive the product at room temperature, please rest assured, the Capsid Bio Quality Inspection Department has conducted experiments to verify that the normal temperature placement of one month will not affect the biological activity of powder products. After collecting, please store the product according to the requirements described in the datasheet. Most Capsid Bio products are stable under the recommended conditions.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

EU Headquarters
Gajeva ulica 28, HR-10000 Zagreb, Croatia

US Headquarters
12222 Merit Dr STE 130 Dallas, Texas, 75251

UK Headquarters
121 ABILITY HOUSE, BROOKER ROAD, EN9 1JH, ESSEX

Orders: Sales@capsidbio.com
Enquiries: info@capsidbio.com
www.capsidbio.com