Periplocin

Catalog No.C14028 Batch:C1402803

Product Datasheet

Technical Data

Formula	$C_{36}H_{56}O_{13}$					
Molecular Weight	696.82	**CAS No.**	13137-64-9			
Solubility (25°C)*	In vitro	DMSO	100 mg/mL (143.5 mM)			
		Water	Insoluble			
	In vivo (Add solvents to the product individually and in order)	Oral Administration	Homogeneous suspension	CMC-NA	≥5mg/ml	Taking the 1 mL working solution as an example, add 5 mg of this product to 1 ml of CMC-Na solution, mix evenly to obtain a homogeneous suspension with a final concentration of 5 mg/ml.
		Injection	Clear solution	5%DMSO [1]→ 40%PEG300 [2]→ 5%Tween80 [3]→ 50%ddH2O Validated by Capsid Bio labs. Should you need adjustments to this formulation, contact our sales team for custom testing.	3.000mg/ml (4.31mM)	Taking the 1 mL working solution as an example, add 50 μL of 60 mg/ml clarified DMSO stock solution to 400 μL of PEG300, mix evenly to clarify it; add 50 μL of Tween80 to the above system, mix evenly to clarify; then continue to add 500 μL of ddH2O to adjust the volume to 1 mL. The mixed solution should be used immediately for optimal results.

* <1 mg/ml means slightly soluble or insoluble.
* Please note that Capsid Bio tests the solubility of all compounds in-house, and the actual solubility may differ slightly from published values. This is normal and is due to slight batch-to-batch variations.
* Room temperature shipping (Stability testing shows this product can be shipped at 2-8°C for storage purposes.)

Preparing Stock Solutions

Concentration \ Volume / Mass	1 mg	5 mg	10 mg
1 mM	1.4351 mL	7.1755 mL	14.3509 mL
5 mM	0.2870 mL	1.4351 mL	2.8702 mL
10 mM	0.1435 mL	0.7175 mL	1.4351 mL
50 mM	0.0287 mL	0.1435 mL	0.2870 mL

Biological Activity

Description	Periplocin (Periplocoside), extracted from the traditional herbal medicine cortex periplocae, has cardiac and anticancer activity. This compound could significantly boost proliferation, migration and stimulate collagen production in fibroblast L929 cells, which is dependent on activation of Src/ERK and PI3K/Akt pathways mediated by **Na/K-ATPase**, and thus promoting wound healing.

RETURN POLICY

Capsid Bio's Unconditional Return Policy ensures a smooth online shopping experience for our customers. If you are in any way unsatisfied with your purchase, you may return any item(s) within 7 days of receiving it. In the event of product quality issues, either protocol related or product related problems, you may return any item(s) within 365 days from the original purchase date. Please follow the instructions below when returning products.

SHIPPING & STORAGE

Capsid Bio products are transported at room temperature. If you receive the product at room temperature, please rest assured, the Capsid Bio Quality Inspection Department has conducted experiments to verify that the normal temperature placement of one month will not affect the biological activity of powder products. After collecting, please store the product according to the requirements described in the datasheet. Most Capsid Bio products are stable under the recommended conditions.

NOT FOR HUMAN, VETERINARY DIAGNOSTIC OR THERAPEUTIC USE.

EU Headquarters
Gajeva ulica 28, HR-10000 Zagreb, Croatia

US Headquarters
12222 Merit Dr STE 130 Dallas, Texas, 75251

UK Headquarters
121 ABILITY HOUSE, BROOKER ROAD, EN9 1JH, ESSEX

Orders: Sales@capsidbio.com
Enquiries: info@capsidbio.com
www.capsidbio.com