Phospho-SEK1/MKK4 (Ser257) Antibody [C11A7]

Catalog No.C05290 Batch:C0529003

Product Datasheet

Biological Description

Specificity	Phospho-SEK1/MKK4 (Ser257) Antibody [C11A7] detects endogenous levels of SEK1/MKK4 phosphorylated at Ser257 .
Background	SEK1, or alternatively known as MKK4 or Jun kinase kinase (JNKK), is a kinase responsible for activating the MAP kinase counterparts SAPK and JNK in response to diverse cellular stresses and inflammatory cytokines. Its activation mechanism involves phosphorylation by MEKK at serine 257 and threonine 261 residues. Similar to MEK, SEK1 exhibits dual-specificity protein kinase activity and targets SAPK/JNK by phosphorylating a conserved TPY site within its activation loop.

Usage Information

Application	WB, FCM	Dilution	WB: 1:1000; FCM: 1:400
Reactivity	Human, Mouse, Rat, Monkey		
Source	Rabbit Monoclonal Antibody	MW	44 kDa
Storage Buffer	PBS, pH 7.2+50% Glycerol+0.05% BSA+0.01% NaN_3		
Storage (from the date of receipt)	–20°C (avoid freeze-thaw cycles), 2 years		

Experimental Methods

WB

Experimental Protocol:

Sample preparation
1. Tissue: Lyse the tissue sample by adding an appropriate volume of ice-cold **RIPA/Nuclear Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**，and homogenize the tissue at a low temperature or lyse it by sonication on ice, then incubate on ice for 30 minutes.
2. Adherent cell: Aspirate the culture medium and transfer the cells into an EP tube. Wash the cells with ice-cold PBS twice. Add an appropriate volume of **RIPA/Nuclear Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**, sonicate to lyse the cells, and incubate on ice for 30 minutes.
3. Suspension cell: Transfer the culture medium to a pre-cooled centrifuge tube. Centrifuge and aspirate the supernatant. Wash the cells with ice-cold PBS twice.Add an appropriate volume of **RIPA/Nuclear Lysis Buffer (containing Protease Inhibitor Cocktail, Phosphatase Inhibitor Cocktail)**, sonicate to lyse the cells, and incubate on ice for 30 minutes.
4. Place the lysate into a pre-cooled microcentrifuge tube. Centrifuge at 4°C for 15 min. Collect the supernatant;
5. Remove a small volume of lysate to determine the protein concentration;
6. Combine the lysate with protein loading buffer. Boil 20 µL sample under 95-100°C for 5 min. Centrifuge for 5 min after cool down on ice.

Electrophoretic separation
1. According to the concentration of extracted protein, load appropriate amount of protein sample and marker onto SDS-PAGE gels for electrophoresis. Recommended separating gel (lower gel) concentration: 10%. Reference Table for Selecting SDS-PAGE Separation Gel Concentrations
2. Power up 80V for 30 minutes. Then the power supply is adjusted (110 V~150 V), the Marker is observed, and the electrophoresis can be stopped when the indicator band of the predyed protein Marker where the protein is located is properly separated. (Note that the current should not be too large when electrophoresis, too large current (more than 150 mA) will cause the temperature to rise, affecting the result of running glue. If high currents cannot be avoided, an ice bath can be used to cool the bath.)

Transfer membrane
1. Take out the converter, soak the clip and consumables in the pre-cooled converter;
2. Activate PVDF membrane with methanol for 1 min and rinse with transfer buffer;
3. Install it in the order of "black edge of clip - sponge - filter paper - filter paper - glue -PVDF membrane - filter paper - filter paper - sponge - white edge of clip";
4. The protein was electrotransferred to PVDF membrane. (0.45 µm PVDF membrane is recommended) Reference Table for Selecting PVDF Membrane Pore Size Specifications
Recommended conditions for wet transfer: **200 mA, 120 min**.
(Note that the transfer conditions can be adjusted according to the protein size. For high-molecular-weight proteins, a higher current and longer transfer time are recommended. However, ensure that the transfer tank remains at a low temperature to prevent gel melting.)

Block
1. After electrotransfer, wash the film with TBST at room temperature for 5 minutes;
2. Incubate the film in the blocking solution **(recommending 5% BSA solution)** for 1 hour at room temperature;
3. Wash the film with TBST for 3 times, 5 minutes each time.

Antibody incubation
1. Use 5% skim milk powder to prepare the primary antibody working liquid (recommended dilution ratio for primary antibody 1:1000), gently shake and incubate with the film at 4°C overnight;
2. Wash the film with TBST 3 times, 5 minutes each time;
3. Add the secondary antibody to the blocking solution and incubate with the film gently at room temperature for 1 hour;
4. After incubation, wash the film with TBST 3 times for 5 minutes each time.

Antibody staining
367. Add the prepared ECL luminescent substrate (or select other color developing substrate according to the second antibody) and mix evenly;
2. Incubate with the film for 1 minute, remove excess substrate (keep the film moist), wrap with plastic film, and expose in the imaging system.

IF

Experimental Protocol：

Specimen Preparation
1. Aspirate liquid, then cover cells to a depth of 2–3 mm with 4% Paraformaldehyde diluted in 1X PBS.
NOTE: Paraformaldehyde is toxic, use only in a fume hood.
2. Fix cells for 15 min at room temperature.
3. Aspirate fixative, rinse three times in 1X PBS for 5 min each.
4. Proceed with Immunostaining.

Immunostaining
1. Add theblocking buffer and incubate for 60 min at RT.
2. Prepare primary antibody diluent in antibody dilution buffer as recommended .
3. Aspirate blocking solution, apply diluted primary antibody.
4. Incubate overnight at 4°C.

Experimental Methods

	5. Rinse three times in 1X PBS for 5 min each. 6. Incubate specimens in fluorochrome-conjugated secondary antibody diluted in antibody dilution buffer for 1–2 hr at room temperature in the dark. 7. Rinse three times in 1X PBS for 5 min each. 8. Mount slides usingmounting medium with DAPI and cover with coverslips. 9. For best results, allow mountant to cure overnight at room temperature. For long-term storage, store slides flat at 4°C protected from light.
IHC	Experimental Protocol： Deparaffinization/Rehydration 1. Deparaffinize/hydrate sections: 2. Incubate sections in three washes of xylene for 5 min each. 3. Incubate sections in two washes of 100% ethanol for 10 min each. 4. Incubate sections in two washes of 95% ethanol for 10 min each. 5. Wash sections two times in dH2O for 5 min each. 6.Antigen retrieval: For Citrate: Heat slides in a microwave submersed in 1X citrate unmasking solution until boiling is initiated; continue with 10 min at a sub-boiling temperature (95°-98°C). Cool slides on bench top for 30 min. Staining 1. Wash sections in dH2O three times for 5 min each. 2. Incubate sections in 3% hydrogen peroxide for 10 min. 3. Wash sections in dH2O two times for 5 min each. 4. Wash sections in wash buffer for 5 min. 5. Block each section with 100–400 µl of blocking solution for 1 hr at room temperature. 6. Remove blocking solution and add 100–400 µl primary antibody diluent in to each section. Incubate overnight at 4°C. 7. Remove antibody solution and wash sections with wash buffer three times for 5 min each. 8. Cover section with 1–3 drops HRPas needed. Incubate in a humidified chamber for 30 min at room temperature. 9. Wash sections three times with wash buffer for 5 min each. 10. Add DAB Chromogen Concentrate to DAB Diluent and mix well before use. 11. Apply 100–400 µl DAB to each section and monitor closely. 1–10 min generally provides an acceptable staining intensity. 12. Immerse slides in dH2O. 13. If desired, counterstain sections with hematoxylin. 14. Wash sections in dH2O two times for 5 min each. 15. Dehydrate sections: Incubate sections in 95% ethanol two times for 10 sec each; Repeat in 100% ethanol, incubating sections two times for 10 sec each; Repeat in xylene, incubating sections two times for 10 sec each. 16. Mount sections with coverslips and mounting medium.

References

https://pubmed.ncbi.nlm.nih.gov/7855889/ [2] Kyriakis JM, et al. Nature. 1994 May 12;369(6476):156-60.

Application Data

WB

Validated by Capsid Bio



Lane 1: Hela
Lane 2: Hela (sorbitol treated)
Lane 3: NIH/3T3
Lane 4: NIH/3T3 (UV treated)